


Pythagoras' theorem

Introduction

Pythagoras' theorem relates the lengths of the sides of a right-angled triangle. This leaflet reminds you of the theorem and provides some revision examples and exercises.

1. Pythagoras' theorem

Study the right-angled triangle shown.




In any right-angled triangle, ABC , the side opposite the right-angle is called the **hypotenuse**. Here we use the convention that the side opposite angle A is labelled a . The side opposite B is labelled b and the side opposite C is labelled c .

Pythagoras' theorem states that the square of the hypotenuse, (c^2) , is equal to the sum of the squares of the other two sides, $(a^2 + b^2)$.

$$\text{Pythagoras' theorem: } c^2 = a^2 + b^2$$

Example



Suppose $AC = 9\text{cm}$ and $BC = 5\text{cm}$ as shown. Find the length of the hypotenuse, AB .

Solution


Here, $a = BC = 5$, and $b = AC = 9$. Using the theorem

$$\begin{aligned}c^2 &= a^2 + b^2 \\ &= 5^2 + 9^2 \\ &= 25 + 81 \\ &= 106 \\ c &= \sqrt{106} = 10.30 \quad (2\text{dp.})\end{aligned}$$

The hypotenuse has length 10.30cm.

Example

In triangle ABC shown, suppose that the length of the hypotenuse is 14cm and that $a = BC = 3$ cm. Find the length of AC .



Solution

Here $a = BC = 3$, and $c = AB = 14$. Using the theorem

$$\begin{aligned}c^2 &= a^2 + b^2 \\ 14^2 &= 3^2 + b^2 \\ 196 &= 9 + b^2 \\ b^2 &= 196 - 9 \\ &= 187 \\ b &= \sqrt{187} = 13.67 \quad (2\text{dp.})\end{aligned}$$

The length of AC is 13.67cm.

Exercises

1. In triangle ABC in which $C = 90^\circ$, $AB = 25$ cm and $AC = 17$ cm. Find the length BC .
2. In triangle ABC , the angle at B is the right-angle. If $AB = BC = 5$ cm find AC .
3. In triangle CDE the right-angle is E . If $CD = 55$ cm and $DE = 37$ cm find EC .

Answers

1. 18.33 cm. (2dp.)
2. $AC = \sqrt{50} = 7.07$ cm. (2dp.)
3. $EC = \sqrt{1656} = 40.69$ cm. (2dp.)