

## Product and quotient rules

### Introduction

As their names suggest, the **product rule** and the **quotient rule** are used to differentiate products of functions and quotients of functions. This leaflet explains how.

### 1. The product rule

It is appropriate to use this rule when you want to differentiate two functions which are multiplied together. For example

$$y = e^x \sin x \quad \text{is a product of the functions } e^x \text{ and } \sin x$$

In the rule which follows we let  $u$  stand for the first of the functions and  $v$  stand for the second.

If  $u$  and  $v$  are functions of  $x$ , then

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

### Example

If  $y = 7xe^{2x}$  find  $\frac{dy}{dx}$ .

### Solution

Comparing the given function with the product rule we let

$$u = 7x, \quad v = e^{2x}$$

It follows that

$$\frac{du}{dx} = 7, \quad \text{and} \quad \frac{dv}{dx} = 2e^{2x}$$

Thus, using the product rule,

$$\frac{d}{dx}(7xe^{2x}) = 7x(2e^{2x}) + e^{2x}(7) = 7e^{2x}(2x + 1)$$

## 2. The quotient rule

It is appropriate to use this rule when you want to differentiate a quotient of two functions, that is, one function divided by another. For example

$$y = \frac{e^x}{\sin x} \quad \text{is a quotient of the functions } e^x \text{ and } \sin x$$

In the rule which follows we let  $u$  stand for the function in the numerator and  $v$  stand for the function in the denominator.

If  $u$  and  $v$  are functions of  $x$ , then

$$\frac{d}{dx} \left( \frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

### Example

If  $y = \frac{\sin x}{3x^2}$  find  $\frac{dy}{dx}$ .

### Solution

Comparing the given function with the quotient rule we let

$$u = \sin x, \quad \text{and} \quad v = 3x^2$$

It follows that

$$\frac{du}{dx} = \cos x \quad \text{and} \quad \frac{dv}{dx} = 6x$$

Applying the quotient rule gives

$$\frac{dy}{dx} = \frac{3x^2 \cos x - \sin x (6x)}{9x^4} = \frac{3x(x \cos x - 2 \sin x)}{9x^4} = \frac{x \cos x - 2 \sin x}{3x^3}$$

### Exercises

Choose an appropriate rule in each case to find  $\frac{dy}{dx}$ .

1.  $y = x^2 \sin x$

2.  $y = e^x \cos x$

3.  $y = \frac{e^x}{x^2+1}$

4.  $y = \frac{x^2+1}{e^x}$

5.  $y = 7x \log_e x$

6.  $y = \frac{x-1}{\sin 2x}$

### Answers

1.  $x^2 \cos x + 2x \sin x$     2.  $-e^x \sin x + e^x \cos x = e^x(\cos x - \sin x)$     3.  $\frac{e^x(x^2-2x+1)}{(x^2+1)^2}$

4.  $\frac{2x-x^2-1}{e^x}$ ,    5.  $7(1 + \log_e x)$ ,    6.  $\frac{\sin 2x - 2(x-1) \cos 2x}{\sin^2 2x}$ .